



Alliance

(<https://www.aquaculturealliance.org>)



**Global
Aquaculture
Advocate**[™]

Aquafeeds

A comparison of U.S. shrimp grow-out diets

Sunday, 1 December 2002

By Leonard G. Obaldo, Ph.D.

O.I. begins comparative testing





Commercial U.S.-made shrimp feeds provided good survival but higher FCR when compared to an expensive control feed.

Although the nutritional and physicochemical composition of shrimp feeds plays a vital role in establishing diet quality indices for optimum animal performance, there is limited information on the quality characteristics and performance of shrimp grow-out diets manufactured in the United States. Databases on diet characteristics could be generated to provide baseline information and a reference for improving U.S. shrimp diets, and help identify priority areas for feed research.

In 2001, with support from a USDA-ARS grant, the Aquatic Feeds and Nutrition program of the Oceanic Institute (O.I.) in Hawaii, USA evaluated commercially available shrimp growout diets manufactured in the U.S. The research studied the nutrient composition, pellet water stability, leaching characteristics, and biological performance of the diets.

Sample collection

At the Oceanic Institute, samples of shrimp growout diets were collected from major U.S. feed manufacturers and suppliers. Out of seven companies contacted, only four responded, with a total of 15 diets received for evaluation. The diets had a pellet diameter of 2.4 mm, protein content ranging 27 to 56 percent, and fat content of 6 to 12 percent on a dry matter basis.

Pellet analysis

Together with two control feeds (an O.I. experimental diet and Japanese commercial shrimp diet), the U.S. diets were analyzed for proximate composition, pellet water stability, and pellet leaching. Diet proximate composition was determined using O.I. standard laboratory procedures. Pellet water stability and leaching were determined by immersing 2 grams of samples for two hours in saltwater at 25 degrees-C temperature with no pellet agitation and water movement.

Growth trial

A four-week growth trial was conducted to evaluate the biological performance of the diets. Shrimp of about 1.38 grams were stocked in 52-liter rectangular glass aquaria using three aquaria per dietary treatment. Initial stocking density was 12 shrimp per aquarium (50 shrimp per square meter). A seawater flow-through system with 100 percent hourly water exchange and an aeration rate of three liters of air per minute was maintained throughout the trial.

All treatments were fed with a fixed ration by automatic feeders eight times daily. Shrimp were weighed at the start and end of the growth trial and evaluated for survival, final body weight, body weight increase and feed-conversion ratio (FCR).

Results

Of the 15 diets tested, the best-performing diet from each U.S. feed company (identified as companies A, B, C and D to protect anonymity) was selected and compared with the Japanese control diet. Although very expensive (about U.S. \$3.00 per kg), this control diet was used because previous growth trials at O.I. found it the best-performing commercial shrimp grow-out diet with good attractability and water stability.

Physicochemical characteristics

The best U.S. diets had much lower protein contents, higher fat contents, and lower ash contents than the Japanese control diet (Table 1). The percentages of dry matter, protein, fat and ash remaining after two hours of leaching in saltwater are presented in Table 2.

Obaldo, Proximate composition (dry matter basis) of best-performing diets, Table 1

Diet	Protein (%)	Fat (%)	Ash (%)	Dry Matter (%)
Company A	46.0	11.4	11.7	91.6
Company B	42.4	7.5	10.0	92.0
Company C	47.1	10.7	10.4	91.1
Company D	44.9	9.2	9.1	92.2
Control (O.I. 40)	42.4	10.7	6.1	91.6
Control (Japan)	66.2	6.9	16.0	91.9

Table 1. Proximate composition (dry matter basis) of best-performing diets from U.S. feed companies.

Obaldo, Composition of best-performing U.S. diets, Table 2

Diet	Remaining Protein (%)	Remaining Fat (%)	Remaining Ash (%)	Remaining Dry Matter (%)
Company A	92	92	74	93
Company B	93	80	92	91
Company C	87	81	88	88
Company D	89	92	91	90
Control (O.I. 40)	80	78	88	86
Control (Japan)	92	57	72	89

Table 2. Composition of best-performing U.S. diets after 2 hours of leaching in saltwater.

The data showed little variation in the results for percent protein and percent dry matter remaining, but indicated a significant difference on percent fat remaining after leaching. The Japanese control diet had 57 percent percent fat remaining, as compared to 80 to 92 percent for the U.S. shrimp diets.

Growth performance

In general, the results indicated the best shrimp grow-out diets produced in the U.S. had very good survival but relatively lower growth and higher FCR than the Japanese commercial control diet (Table 3). Although these differences (survival, growth, FCR) were not statistically significant, it is important to note that the samples tested in the study were diets available at the time of request, and did not necessarily represent the best diets produced by each U.S. feed company.

Obaldo, Biological performance of best-performing U.S. diets, Table 3

Diet	Weight (g)	Growth (g/week)	FCR	Survival (%)
Company A	3.35	0.49	1.63	100
Company B	3.33	0.49	1.65	100
Company C	3.45	0.52	1.56	100
Company D	3.09	0.42	1.95	100
Control (O.I. 40)	3.20	0.46	1.81	100
Control (Japan)	3.68	0.57	1.49	94

Table 3. Biological performance of best-performing U.S. diets in a four-week growth trial.

The best U.S. diets had lower protein contents, higher fat contents and lower ash contents than the Japanese control diet.

In addition, due to limited sample quantity, the comparison was based on a short indoor growth trial with no additional food from natural productivity such as that found in zero-exchange systems. Evaluation of the diets in a zero water-exchange system could yield different growth results.

Conclusion

Without knowledge of the ingredients and processing used in making the feeds, it is difficult to establish a consistent relationship between diet nutritional quality and shrimp biological performance. However, there is a need to continue the formulation and development of improved diets to address the requirements of evolving production systems.

Further work is recommended to evaluate and compare the performance of U.S. shrimp grow-out diets in zero-exchange systems. The information generated provides a reference for future research on shrimp feed development.

(Editor's Note: This article was originally published in the December 2002 print edition of the Global Aquaculture Advocate.)

Author



LEONARD G. OBALDO, PH.D.

Oceanic Institute
41-202 Kalanianaʻole Highway
Waimanalo, Hawaii 96795 USA

lobaldo@oceanicinstitute.org (<mailto:lobaldo@oceanicinstitute.org>)

Copyright © 2016–2020 Global Aquaculture Alliance

All rights reserved.