

**Alliance**<https://www.aquaculturealliance.org>

Aquafeeds

# Cassia seeds a potential protein source in tilapia feed

Friday, 1 August 2003

By L.C. Nwana, O.A. Fagbenro and T. Jegede

## Seedmeal can serve as a good source of minerals in fish diets

Conventional plant protein ingredients used in fish nutrition – such as soybean meal and groundnut cake – can be scarce and expensive due to competition for their use in both human and livestock feeds. In 2002, the first author reported that some underexploited forest seeds with high protein levels and suitable amino acid profiles could be used as replacements for expensive plant protein feedstuffs in practical diets for fish.




Fig. 1. ADC protein in tilapia fed cassia meal.

## Study setup

A study by the authors evaluated the nutritive value of cassia (*Cassia fistula*), an underutilized forest seed, and its digestibility in Nile tilapia (*Oreochromis niloticus*), a prominent freshwater aquaculture species.

Cassia seed bunches were harvested from the campus of the Federal University of Technology in Akure, Nigeria. The seeds were removed and roasted in a frying pan at 70 to 90 degrees-C for 15 to 20 minutes to reduce the moisture content and substantially reduce the anti-nutrients in cassia seeds. The seeds were cleaned and milled into fine powder to form a meal that was analyzed for proximate and mineral compositions according to standard methods.

In digestibility trials, the cassia seed meal replaced soybean meal at 0, 25, 50, 75 and 100 percent in 35 percent-crude protein diets. The diets were fed to triplicate groups of 10-g tilapia in glass tanks at 5 percent body weight per day for 14 days. Fecal samples were collected eight hours after each feeding, pooled and analyzed for proximate composition. Digestibility was calculated using the acid in-soluble ash methods.




Fig. 2. ADC energy in tilapia fed cassia meal.

## Results

Table 1 presents the proximate composition of the *Cassia fistula* meal. Its protein content was high (26 percent), with low ash and fat contents, indicating that meal from cassia seeds could be used in fish nutrition. This protein level was similar to the respective values of 24.5 percent and 25.2 percent obtained by Garg et al (2002) for moong and cowpea seeds used as ingredients in tilapia diets.

The mineral composition of the cassia seed meal (Table 2) showed high values close to those in warmwater fish, indicating the meal can serve as a good source of minerals in fish diets. This composition compared well with the mineral contents of *Cassia laevigata* reported by Siddhuraju et al (1995).

Fish fed a control diet without cassia meal had the highest apparent protein digestibility (ADC protein), which was similar to the ADC protein of the fish fed diets with 25 percent cassia meal (Fig. 1). However, these ADC values were significantly higher than those for the fish fed diets with 50 percent, 75 percent, and 100 percent cassia meal. Also, the ADC values decreased continually as the level of cassia meal in the diets increased. The ADC energy (Fig. 2) of the fish fed diets of 0 percent, 25 percent, 50 percent, and 75 percent of cassia meal were similar, but differed significantly from the value for fish fed 100v cassia meal.

## Nwanna, Proximate composition of *Cassia fistula* flour, Table 1

Nutrient	Percentage
Crude protein	26.0
Crude fat	3.14
Crude fiber	8.05
Ash	2.49
Moisture	9.82
Nitrogen free extract	50.5

Table 1. Proximate composition of *Cassia fistula* flour.

## Nwanna, Mineral composition of *Cassia fistula* flour, Table 2

Mineral	b/kg
Phosphorous	11.2
Potassium	3.7
Magnesium	2.0
Calcium	1.6
Sodium	1.2
Iron	0.3
Zinc	0.13

Table 2. Mineral composition of *Cassia fistula* flour.

Based on these results, soybean meal could be replaced by cassia meal at a 25 percent inclusion level in *O. niloticus* diets for fish in the tester size range.

*Note: Cited references are available from the first author.*

*(Editor's Note: This article was originally published in the August 2003 print edition of the Global Aquaculture Advocate.)*

## Authors

---



**L.C. NWANNA**

Department of Fisheries and Wildlife  
Federal University of Technology  
PMB 704 Akure, Nigeria

[drlu2001@yahoo.com](mailto:drlu2001@yahoo.com) (<mailto:drlu2001@yahoo.com>).



**O.A. FAGBENRO**

Department of Fisheries and Wildlife  
Federal University of Technology  
PMB 704 Akure, Nigeria



**T. JEGEDE**

Department of Fisheries and Wildlife  
Federal University of Technology  
PMB 704 Akure, Nigeria

Copyright © 2016–2020 Global Aquaculture Alliance

All rights reserved.