



ALLIANCE™

(<https://www.globalseafood.org>).



 Responsibility

Correct liming improves pond water, bottom quality

1 August 2002

By Claude E. Boyd, Ph.D.

Chalk, marl, crushed seashells and wood ash are not as effective as calcite



Shrimp farmers often apply liming materials during pond preparation to improve the pH of bottom soils.

Shrimp and fish farmers frequently apply liming materials to ponds to increase the pH of bottom soils and elevate alkalinity and hardness in water. These chemical changes improve conditions for microbial activity and benthic animals; increase the availability of carbon dioxide, phosphorus, and other nutrients; enhance phytoplankton growth; and improve the survival and growth of the aquaculture crop.

Confusion about the properties of liming materials, however, often results in these materials being used incorrectly. And, of course, if soil or water are not acidic, treatment with agricultural limestone will be of little or no benefit.

Common forms

The most common liming materials are agricultural limestone, burnt lime or quick lime, and hydrated lime or slaked lime. All of these products are made from limestone. Limestone is a relatively soft rock composed of calcium carbonate (CaCO_3 or calcite), calcium magnesium carbonate ($\text{CaCO}_3 \cdot \text{MgCO}_3$ or dolomite), or a mixture of these carbonates.

Solubility



(<https://globalseafood.typeform.com/podcastq124>).

Like most rocks, limestone in raw form is not highly soluble. By pulverizing it into fine particles, the surface area:mass ratio is increased to enhance solubility, as measured by increase in alkalinity (Fig. 1). In good-quality agricultural limestone, most particles will pass a 60-mesh sieve (0.25-mm openings). In high-quality agricultural limestone, most particles will pass a 100-mesh sieve (0.15-mm openings).




Fig. 1: Alkalinity of pulverized agricultural limestone.

Color

The color of agricultural limestone is variable. Most samples are a shade of white, but can also appear red, yellow, or brown.

Other liming materials

Other materials that have been used for liming include chalk (soft limestone), marl (loose carbonate deposits often mixed with clay), crushed seashells, and wood ash. Most are not as effective as calcite, and some of their byproducts can contain harmful levels of heavy metals or other undesirable contaminants.

Chemical reactions

When agricultural limestone reacts with acidity, the acidity is neutralized and calcium carbonate is transformed to calcium ions, carbon dioxide, and water. Magnesium carbonate in agricultural limestone reacts with acidity in a similar manner, but magnesium ions instead of calcium ions are released.

The most important reaction of agricultural limestone in water is with carbon dioxide. This reaction results in dissolution of the calcium carbonate and an increase in the calcium and bicarbonate concentrations in solution. Magnesium carbonate reacts in a similar manner, but magnesium and bicarbonate result.

Water hardness

The total hardness of water is a measure of the calcium and magnesium concentration expressed as equivalent calcium carbonate. For example, suppose a water contains 5 milligrams per liter calcium and 1 milligrams per liter magnesium. The total hardness in equivalent calcium carbonate can be determined as:

$$5 \text{ mg Ca/l} \times \frac{100 \text{ mg CaCO}_3/\text{mM}}{40 \text{ mg Ca/mM}} = 12.5 \text{ mg/l CaCO}_3/\text{l}$$

$$1 \text{ mg Mg/l} \times \frac{100 \text{ mg CaCO}_3/\text{mM}}{24.31 \text{ mg Mg/mM}} = 4.12 \text{ mg CaCO}_3/\text{l}$$

$$\text{Total hardness} = 16.62 \text{ mg CaCO}_3/\text{l}$$

In practice, total hardness can be measured directly by titration of calcium and magnesium with ethylenediamine-tetraacetic acid (EDTA).

Alkalinity

The total alkalinity of water results from the titratable bases expressed in terms of equivalent calcium carbonate. Bicarbonate usually is the main base in water. Alkalinity is measured by titration of a water sample to pH 4.5 with standard acid and estimation of the calcium carbonate equivalence of the amount of acid used.

Limestone and water

Application of agricultural limestone to water normally increases total alkalinity and hardness by roughly equal amounts. However, the solubility of calcium carbonate in water in equilibrium with normal atmospheric concentrations of carbon dioxide is about 60 milligrams per liter. In ponds, there usually is more carbon dioxide available from organic matter decomposition, and calcium carbonate is more soluble. Nevertheless, if pond waters contain more than 80 to 100 milligrams per liter alkalinity, agricultural limestone usually will not dissolve.

Limestone and soil

Agricultural limestone reacts in soil much the same way it does in water. However, the calcium and magnesium ions that result from its reaction with hydrogen ions or carbon dioxide exchange for aluminum and iron ions absorbed on soil colloids. This reduces the acidity of the soil and causes its pH to increase. The pH of soil cannot be increased above 8.3 by application of agricultural limestone.

Liming effects

The primary effects of agricultural limestone applications in aquaculture systems are to decrease acidity and increase the pH, hardness, and alkalinity of acidic soils and water. These chemical changes improve conditions for microbial activity and benthic animals; increase the availability of carbon dioxide, phosphorus, and other nutrients; enhance phytoplankton growth; and improve the survival and growth of the aquaculture crop. Of course, if soil or water are not acidic, treatment with agricultural limestone will be of little or no benefit.

Burnt and hydrated lime

Burnt lime is made by burning limestone in a furnace at high temperature. Hydrated lime is made by treating burnt lime with water. Both burnt lime and hydrated lime usually are fine, white powders. When put into ponds, calcium oxide reacts with water and forms calcium hydroxide, which neutralizes acidity and causes pH to rise. Calcium hydroxide also can react with carbon dioxide.

When added to water, burnt lime and hydrated lime have the same effects as agricultural limestone, but initially cause a much higher pH. Burnt and hydrated limes should not normally be applied to pond water during the culture period to prevent potentially harmful high pH levels. These two materials, however, can be applied to the bottoms of empty ponds at 1,000 to 2,000 kilograms per hectare to raise pH and kill pathogens and other unwanted organisms.

Neutralizing values

The strength of liming materials is reported as a neutralizing value, with the value of pure calcium carbonate established as 100 percent. Based on this system, the neutralizing values of pure compounds used as liming material are:

Calcite 100 percent

Dolomite 108 percent

Hydrated Lime 135 percent

Burnt Lime 178 percent

Agricultural limestone made from pure dolomite is 1.08 times as strong in neutralizing acidity as that made from calcite. Burnt lime and hydrated lime are, respectively, 1.78 and 1.35 times as strong in neutralizing acidity as agricultural limestone made from calcite. Of course, individual samples of liming materials will vary because of impurities.

(Editor's Note: This article was originally published in the August 2002 print edition of the Global Aquaculture Advocate.)

Author



CLAUDE E. BOYD, PH.D.

Professor, Department of Fisheries and Allied Aquacultures
International Center for Aquaculture and Aquatic Environments
Auburn University
Auburn, Alabama 36849 USA

boydce1@auburn.edu (<mailto:boydce1@auburn.edu>)

Copyright © 2024 Global Seafood Alliance

All rights reserved.