



ALLIANCE™

(<https://www.globalseafood.org>).



 Aquafeeds

Evaluating brown seaweed supplementation in feed for Atlantic salmon smolts

18 May 2020

By Collins Kamunde, Ph.D. , Ravinder Sappal, DVM, Ph.D. and Tarek Mostafa Melegy

Results indicate increased food intake, improved growth



Results of this study to evaluate the effect of addition of a kelp meal to feed for Atlantic salmon smolts suggest that inclusion levels up to 10 percent can increase feed consumption, enhance growth performance and improve antioxidant capacity and alleviate adverse effects of stressors like temperature. Photo by Nicolás from seno de reloncaví, Chile / CC BY-SA (<https://creativecommons.org/licenses/by-sa/2.0>).

There is continued interest in the use of plant ingredients in aquafeeds, but much remains unknown about the value of aquatic macroalgae (seaweeds) in the nutrition of aquatic animals. The most relevant effects of seaweed supplementation in aquaculture feeds include stimulation of growth performance, enhancement of feed utilization efficiency, improvement of nutrient assimilation, and improvement of fatty acid profile in muscle.

In addition, seaweeds contain a wide array of bioactive compounds and secondary metabolites with potential utility as phytonutrients and nutraceuticals in animal feed. There is much interest in determining if these bioactive compounds can improve the overall health status of fish by enhancing resistance to disease, improving antioxidant capacity, and/or alleviating routine aquaculture stress associated with crowding or events such as handling (e.g., during grading and vaccination) and transportation.

At high inclusion levels, seaweeds have been shown to impair fish growth performance and feed efficiency, so determining the inclusion levels that improve fish growth performance and/or health status remains the primary focus of most relevant studies. Such knowledge could permit the replacement of expensive ingredients of fish feed – such as fishmeal and/or mixing of small amounts of seaweeds or their extracts – with finished aquafeeds to harness the other beneficial effects not related to growth.

However, the variability in the biochemical composition and inconsistent effects of seaweed supplementation among fish necessitates testing the effects of supplementation of specific seaweed on specific fish species. In particular, salmonids are the most important aquaculture fish group and use the greatest volumes of fishmeal and fish oil in aquafeeds; therefore, demonstrating a role of seaweeds in salmonid nutrition would have major implications for aquaculture.



(<https://link.chtbl.com/aquapod>).

This article – adapted and summarized from the [original](https://doi.org/10.1371/journal.pone.0219792) (<https://doi.org/10.1371/journal.pone.0219792>) – assessed the effect of the addition of AquaArom, a commercial meal from brown seaweed or kelp (*Laminaria* sp.), to feed and the growth performance and other parameters in smolts of Atlantic salmon (*Salmo salar*). We are grateful to Nicole MacDonald for technical assistance.

Study setup

The feeding trial was conducted at the aquatic facility of the Atlantic Veterinary College, University of Prince Edward Island (Canada). Atlantic salmon smolts (initial average weight: 77 grams) were obtained from Northern Harvest in Cardigan and were maintained and acclimated for one month in a 1,200-liter tank with flow-through aerated well-water. After the acclimation period, the smolts (average weight 92 grams) were randomly distributed in groups of eight in 160-liter tanks comprising triplicates for each of the four experimental groups (control and 3, 6 and 10 percent seaweed supplementation). The tanks were then randomly distributed in a 2 × 6 block within the experimental room. A photoperiod of 12 hours of light and 12 hours of dark was maintained.

The fish were hand-fed the designated diet twice a day for 30 days. After dispensing the food into the tanks, fish were allowed to feed for 1 hour, following which the uneaten food was collected, dried and subtracted from the total amount of food dispensed to estimate food consumption per tank, feed conversion ratio (FCR) and protein efficiency ratio (PER). Bulk fish weights obtained weekly were used to calculate the ration for the following week. On day 30, fish were euthanized, individually weighed and measured. Blood samples were collected, and the fish dissected to harvest livers and viscera.

The experimental diets were prepared in-house by supplementing finished commercial fish feed – EWOS micro crumble for salmonids (Ewos Canada Ltd, St. George, New Brunswick, Canada) – with the required amount of brown seaweed flakes (AquaArom) prepared from kelp (*Laminaria* sp.) and provided by ADDiCAN Inc., Canada. Diets were calculated to deliver 0 (control), 3 (SW-3), 6 (SW-6), and 10 (SW-10) percent seaweed on a dry matter basis.

For detailed information on the experimental design and diets; analysis of food and seaweed composition; feeding trial and sampling; analyses of biological samples; and statistical analyses, refer to the original publication.

Results and discussion

Analysis of the seaweed showed that it contained 12.37 percent crude protein and non-detectable levels of fat on dry matter basis. Additionally, relative to the control commercial salmonid feed, the seaweed contained lower levels of key minerals (phosphorus, copper, and zinc) and higher levels of neutral detergent fiber (NDF), ash and NaCl. Addition of the seaweed to commercial salmonid feed caused slight changes in the proportion of other dietary ingredients. Notably, the crude protein decreased from 57.6 percent (control) to 52.7 (10 percent seaweed) and from 54.7 to 48.9 percent (as fed basis) in control and 10 percent seaweed-supplemented food, respectively.

To put these values into nutritional requirement context, levels of crude protein in feeds for various age classes of Atlantic salmon range from 42 to 50 percent, with 45 percent for juveniles. Similarly, the slightly reduced levels of phosphorus, copper, and zinc in the seaweed-supplemented diets more than met the respective minimum nutritional requirements for various life stages/size classes of Atlantic salmon. Based on these findings, it is likely that higher rates of inclusion (>10 percent) of this seaweed might decrease levels of key ingredients to levels below their minimum requirements.

There were no mortalities during the 30-day feeding trial; all of the fish in each of the 12 tanks survived and no malformations or other adverse effects were observed. The fish readily accepted the experimental food and the overall effect of seaweed supplementation on food intake (Fig. 1) was significant. Specifically, fish maintained on the seaweed-supplemented diets consumed significantly more food than the control. Consistent with the pattern of food consumption, fish weight per tank varied with seaweed supplementation, with fish maintained on food supplemented with 3 and 10 percent seaweed having significantly higher final body weights than the control.




Fig 1: Brown seaweed supplementation increases food intake and body weight in Atlantic salmon smolts. (A) Daily food intake. (B) Final fish weight. Ctl: control (0 percent seaweed), SW-3: 3 percent seaweed, SW-6: 6 percent seaweed, SW-10: 10 percent seaweed. Bars with different letters are significantly different (one-way ANOVA, LSD test, $p < 0.05$).

The higher final fish weights relative to the control for the 3 and 10 percent seaweed-supplemented fish were associated with higher percent weight gain, higher daily weight gain and higher specific growth rate, SGR (Fig. 2). Although seaweed supplementation overall did not significantly alter the FCR, the 3 percent supplemented group had better FCR than the 6 percent supplemented group. Moreover, PER was significantly altered by seaweed supplementation (Fig. 3) and was higher in the 3 and 10 percent groups, relative to the 6 percent group.

Fig. 2: Effect of brown seaweed supplementation on growth performance indices of Atlantic salmon smolts. (A) percent weight gain. (B) Daily weight gain. (C) SGR. Ctl: control (0 percent seaweed), SW-3: 3 percent seaweed, SW-6: 6 percent seaweed, SW-10: 10 percent seaweed. Bars with different letters are significantly different (one-way ANOVA, LSD test, $p < 0.05$).

Fig 3: Supplementation with brown seaweed improves PER but not FCR in Atlantic salmon smolts. (A) FCR. (B) PER. Ctl: control (0 percent seaweed), SW-3: 3 percent seaweed, SW-6: 6 percent seaweed,

SW-10: 10 percent seaweed. Bars with different letters are significantly different (one-way ANOVA, LSD test, $p < 0.05$).

Seaweed supplementation significantly altered fish standard length in which fish that were fed 3 and 10 percent seaweed-supplemented food were longer than the control at the end of the trial. However, Fulton's condition factor, K (which related length and weight) computed from the final weights and standard lengths remained unchanged (Fig 4). Measurement of other morphometric indices revealed that seaweed supplementation did not alter body depth, hepatosomatic indices (HIS) or viscerosomatic indices (VSI), but it increased weights and lengths of the entire guts and intestines for the 3 and 10 percent supplementation levels relative to the control.

Fig. 4: Effect of brown seaweed supplementation on body length and K of Atlantic salmon smolts. (A) Standard length. (B) Condition factor

(K). Ctl: control (0 percent seaweed), SW-3: 3 percent seaweed, SW-6: 6 percent seaweed, SW-10: 10 percent seaweed. Bars with different letters are significantly different (one-way ANOVA, LSD test, $p < 0.05$).

Although seaweeds are considered natural forage for fish, they remain underutilized as an ingredient in aquafeeds. We tested the potential use of seaweed meal derived from brown seaweed of the genus *Laminaria* (kelp) as an additive to commercial salmonid food using Atlantic salmon smolts. Because of their large size and ease of harvesting, brown seaweeds are generally more amenable to exploitation for animal nutrition than other types of macroalgae. However, brown seaweeds have low nutritional value, containing only 3 to 15 percent crude protein compared with green and red seaweeds that may contain up to 26 and 47 percent crude protein, respectively.

Despite their low protein content, brown seaweeds remain popular as potential animal feed additives because they are rich in bioactive compounds. Indeed, brown seaweeds constitute the majority of seaweed used in terrestrial animal nutrition, but they are the least investigated for application in aquafeeds relative to other seaweed classes.

In our study, Atlantic salmon smolts readily accepted food mixed with 3 to 10 percent brown seaweed. Interestingly, smolts maintained on seaweed-supplemented feed consumed more food than the control. This contrasts the typical finding that inclusion of vegetable protein, particularly in high amounts, in fish diets is associated with reduced palatability and food intake, resulting in decreased growth performance. Regardless, increased intake of seaweed-supplemented fish food has also been previously reported and attributed to the presence in seaweeds of various compounds that attract fish to food, thus enhancing consumption.

Another explanation for the observed increased food intake in our study could be that increased fiber content due to the added seaweed would reduce available dietary energy resulting in increased food intake as a mechanism to compensate for the energy shortfalls. And the addition of seaweed to aquafeeds can improve the texture, integrity and water stability of pellets resulting in higher food consumption.

In our study, supplementation with 3 to 10 percent brown seaweed did not affect survival during the 30-day feeding trial. Importantly, fish fed 3 and 10 percent seaweed-supplemented food exhibited higher daily weight gains, SGR and both final weight and total length than the control, while fish fed 6 percent seaweed maintained growth comparable to the control. Moreover, seaweed supplementation increased PER and imposed a strong tendency to improve FCR. Overall, our findings together with various earlier studies indicate that supplementation of aquafeed with seaweeds at appropriate levels can enhance or not alter growth performance in salmonids.

The enhancement of growth by seaweeds has been attributed to seaweed vitamin and mineral content and/or increased lipid mobilization and assimilation of dietary nutrients. In our study, it is unlikely that phosphorus, copper and zinc contributed to the growth-stimulatory effects because the seaweed we tested contained low levels of these minerals. It has also been postulated that prebiotic activity of seaweed components (e.g., polysaccharides and oligosaccharide) stimulates growth of beneficial bacteria, thereby improving digestion and subsequently growth.

While we did not investigate the mechanisms or causes of the enhanced growth performance, we found that the digestive tracts were heavier and longer in fish fed seaweed-supplemented diets, suggesting a larger surface area for digestion and absorption of nutrients. Longer digestive tracts would increase the

retention time of food allowing more time for digestion of seaweed-supplemented diets. Surprisingly, the highest seaweed inclusion level (10 percent) we tested resulted in better growth performance despite a 5 percent-point lower protein content relative to the control. Regardless, the protein contents for all the seaweed-supplemented diets were within the optimal range for Atlantic salmon nutrition.

Overall, while it appears that up to 10 percent inclusion of AquaArom stimulates growth in Atlantic salmon smolts, a longer feeding trial would be necessary for a more definitive conclusion on growth performance. Furthermore, future studies should investigate the mechanisms and components underlying the growth-enhancing effects of this brown seaweed meal.

Our findings are consistent with previous reports that dietary seaweed supplementation modulates antioxidant status and oxidative stress in farmed animals. Thus our study, together with these earlier reports, suggests that dietary brown seaweed supplementation enhances antioxidant capacity directly by increasing levels of antioxidant compounds in circulation/tissues and indirectly by modulating activities of antioxidant defense systems.

Perspectives

Overall, our study shows that the addition of the seaweed meal to commercial salmonid feed increases food intake and enhances growth performance, improves plasma antioxidant capacity, and alleviates the effect of temperature rise on mitochondrial respiration. The slight decline in crude protein and minerals resulting from the addition of up to 10 percent seaweed meal to aquafeed appear to have no adverse consequences on Atlantic salmon smolts. Thus, mixing of this brown seaweed meal with commercial aquafeeds (and potentially feeds for other farm animals) could offer a cost-effective way of harnessing the beneficial effects of seaweeds in animal production.

Authors



COLLINS KAMUNDE, PH.D.

Corresponding author
Department of Biomedical Sciences, Atlantic Veterinary College
University of Prince Edward Island, Canada

ckamunde@upei.ca (mailto:ckamunde@upei.ca).



RAVINDER SAPPAL, DVM, PH.D.

Department of Biomedical Sciences, Atlantic Veterinary College
University of Prince Edward Island, Canada



TAREK MOSTAFA MELEGY

Department of Biomedical Sciences, Atlantic Veterinary College
University of Prince Edward Island, Canada

Copyright © 2024 Global Seafood Alliance

All rights reserved.