



ALLIANCE™

(<https://www.globalseafood.org>)




 Responsibility

Feed trays or broadcasting?

2 September 2012

By Dr. Carlos A. Ching and Dr. Chalor Limsuwan

Broadcast method reduces organic matter in shrimp ponds



This aerial view of an Ecuadorian shrimp farm using the “all feed in trays” method shows organic matter accumulation on pond bottoms in tray areas. Photo by Sebastian Arias.

Three Latin American shrimp farms visited by one of the authors applied all feed via 20 feed trays/ha to shrimp stocked at 20/m² or more in earthen ponds. Depressions were observed in these ponds bottoms after harvest, and the average amount of feed for each tray was over 1 kg in the last month of each production cycle.

Measurements of organic matter concentrations on pond bottoms showed significant differences between feeding and non-feeding areas (Table 1). Values for feed tray areas were above the recommended 3 percent concentration.

Ching, Average soil organic matter, Table 1

Country	Organic Matter Feed Tray Areas	Organic Matter Outside Tray Areas
Brazil	5.35%	2.50%
Ecuador	3.50%	1.76%
Peru	4.23%	2.05%

Table 1. Average soil organic matter concentrations at three shrimp farms.

To address this, the farms decided to level their pond bottoms and changed their feeding methods to broadcast the feed, using only 3 feed trays/ha for feed samples to evaluate consumption. Since then, soil quality improved and shrimp survival increased.

(<https://bspcertification.org/>).

Intensification

Many shrimp farms using earthen ponds in Latin America have applied the “all feed in trays” method for many years. With low stocking densities and proper pond bottom preparation, high concentrations of organic matter in the soil were avoided. However, in recent years, stocking densities increased at many Latin American farms, and the amount of feed applied in each tray rose to more than 1 kg/tray.

When some of the added feed went underneath the trays, thousands of shrimp dug under the trays and caused depressions. In these depressions, the concentrations of organic matter from feed, feces and exuviae increased above the 3 percent recommended value in every cycle of production and led to bacterial disease outbreaks. Thus, many farms are changing their feeding method to broadcasting.



When too much feed is applied in trays, shrimp activity under the trays forms depressions that accumulate organic matter.

Broadcast feeding

Due to the large amount of feed they use, most farms in Asia broadcast all their feed with only small feed samples on feed trays. Latin America is following suit as its farm densities rise.

The method for broadcasting feed used by most of the shrimp farms in the world consists of broadcasting 94 to 97 percent of the feed dose, leaving 3 to 6 percent of the dose to be divided as samples on feed trays. The criteria used for this method depend on the season and size of the pond.

The number of feed trays is defined by the area of the pond (Table 2). In small ponds with intensive culture, more feed trays per hectare are used than in large ponds over 2 ha used for extensive or semi-intensive culture.

Ching, Recommended number of feed tray, Table 2

Pond Area (ha)	Trays/ha
Below 2	6-10
2-5	4-5
5-10	2-3
Over 10	1-2

Table 2. Recommended number of feed tray based on field results at Latin American farms.

Some farms not willing to change their “all feed in trays” approach are overcoming the problem of using too much feed by increasing the number of daily feed doses and/or installing more feed trays, but these changes increase labor costs and logistics.

(Editor’s Note: This article was originally published in the September/October 2012 print edition of the Global Aquaculture Advocate.)

Authors



DR. CARLOS A. CHING

Technical Assistance Manager
Nicovita – Alicorp SAA
Av. Argentina 4793
Callao, Peru

cchingm@alicorp.com.pe (mailto:cchingm@alicorp.com.pe)



DR. CHALOR LIMSUWAN

Professor, Department of Fishery Biology
Kasetsart University
Bangkok, Thailand

Copyright © 2024 Global Seafood Alliance

All rights reserved.