

**Alliance**

(<https://www.aquaculturealliance.org>)



Responsibility

Water quality standards: Total Ammonia Nitrogen

Wednesday, 1 August 2001

By Claude E. Boyd, Ph.D.

Through its Responsible Aquaculture Program, GAA has recommended a range of water quality standards for shrimp farm effluents. Part of a series, this article addresses Total Ammonia Nitrogen (TAN).

Nitrogen is a key factor regulating productivity of natural and agricultural ecosystems because it is an essential plant nutrient, a major component of protein, and often in short supply. Nitrogen occurs with phosphorus in effluents, and these two nutrients are the major cause of eutrophication of water bodies.

Nitrogen cycle

A complex nitrogen cycle functions on a global basis. Most pathways within the cycle (Fig. 1) can be found in all ecosystems, including aquaculture ponds.




Fig. 1. Nitrogen cycle. Arrows indicate movement of nitrogen in shrimp pond. N₂ = nitrogen gas, NO₃ = nitrate, NO₂ = nitrite, NH₃ = nonionized ammonia, NH₄ = ammonium, SON = soluble organic nitrogen.

Vast amounts of nitrogen exist in the atmosphere in gaseous form. This nitrogen can be converted to nitrate by lightning, fixed in organic matter by some microorganisms, and converted to nitrogen for fertilizers by an industrial process. Large quantities of nitrogen are found in living organisms and dead organic matter.

Organisms excrete nitrogenous wastes, and when bacteria and other microorganisms decompose dead organic matter, they release ammonia nitrogen into the environment. Ammonia nitrogen consists of non-ionized ammonia (NH₃) and ammonium ion (NH₄⁺). Nitrate nitrogen is produced when nitrifying organisms oxidize ammonia nitrogen.

Most plants prefer ammonium ion as a nitrogen source, but can also use nitrate as a nitrogen source. Nitrate can be converted back to nitrogen gas by the activity of denitrifying bacteria.

Use in aquaculture

Nitrogen is applied to aquaculture ponds in fertilizers such as urea, ammonium nitrate, and sodium nitrate to promote phytoplankton growth and enhance the production of the culture species. Feeds are commonly used in aquaculture because they allow greater production of the culture species than is possible with fertilizers alone.

Feeds usually contain 20 to 40 percent protein, and protein has an average nitrogen concentration of 16 percent. About 20 to 25 percent of the nitrogen applied to ponds in fertilizers and feeds is retained in the biomass of the culture species and removed from ponds at harvest (Table 1).

Boyd, Typical inputs and outputs of nitrogen, Table 1

Inputs	Inputs	Outputs	Outputs
Feed and fertilizer	90%	Shrimp harvest	25%
Water supply and nitrogen fixation	10%	Denitrification	12%
		Ammonia volatilization	18%
		Sediment accumulation	10%
		Effluent	35%

Reducing eutrophication

By limiting the total ammonia nitrogen concentration in effluents to a concentration that will not cause toxicity in the mixing zone, the potential for eutrophication is also greatly reduced. Limits on total ammonia nitrogen commonly are established in effluent permits, but concentration limits for total nitrogen seldom are specified. Of course, in some permits, there may be a limit on total nitrogen loads.

The calculation of nitrogen load requires data on both effluent volume and total nitrogen concentration. For example, suppose the effluent volume for a shrimp farm is 20,000 cubic meters per day and the average total nitrogen concentration is 4 milligrams per liter (grams per cubic meters). The total nitrogen load is 20,000 cubic meters per day x 4 grams per cubic meter = 80 kilograms per day.

GAA effluent standards

The Global Aquaculture Alliance effluent standard for total ammonia nitrogen is initially 5 milligrams per liter, with a target of 3 milligrams per liter. Compliance with these standards is easy in semi-intensive aquaculture, where feed inputs are usually less than 30-40 kilograms per hectare per day. Total ammonia nitrogen concentrations seldom rise above 1 milligram per liter and are rarely greater than 2 milligrams per liter. Semi-intensive farm managers strive to maintain concentrations below 1 milligram per liter.

Because of the difficulties associated with setting different standards for different production levels, GAA adopted a total ammonia nitrogen standard based on intensive shrimp culture, where concentrations often exceed 5 milligrams per liter and can even reach 10 milligrams per liter. Thus, compliance to GAA's standard will be more difficult in intensive aquaculture.

Reducing ammonia in effluents

Methods for reducing total ammonia nitrogen concentrations in aquaculture pond effluents include the following:

- Use fertilizers in moderate amounts and only when needed to promote phytoplankton growth.
- Do not stock and feed at rates so high that water quality deteriorates in ponds.
- Avoid feeding more than culture species can consume.
- Purchase feeds that contain no more nitrogen than needed by the culture species.
- Use mechanical aeration in intensive ponds to avoid dissolved oxygen concentrations below 3-4 milligrams per liter.

Concentration changes in ponds

Total ammonia nitrogen concentrations in ponds change gradually over a period of days. They tend to increase as culture periods progress and feeding rates increase. Nonionized ammonia concentrations change daily as pH and water temperature increase and decrease in response to inputs of solar radiation.

The analytical methods generally used for ammonia nitrogen do not distinguish between non-ionized ammonia and ammonium. The non-ionized ammonia concentration must be estimated with a table that relates the percentage of non-ionized ammonia to temperature and pH. Such tables can be found in books on water quality in aquaculture.

It is sufficient to measure only total ammonia nitrogen in effluents to comply with GAA standards. Some effluent permits, however, require data on non-ionized ammonia.

Sampling and analysis

Samples for total ammonia nitrogen analyses should be placed in plastic bottles, kept on ice, and analyzed six to eight hours after collection. A private or university laboratory can be contracted to do the analysis, but it often will not be possible to send samples within the required time period. Thus, farm technicians can make the analyses.

For seawater or brackish water, the salicylate method is best for total ammonia nitrogen analysis. The Nessler's method and the phenate procedure are best suited for freshwater. The Hach Chemical Co. of Loveland, Colorado, USA sells total ammonia nitrogen test procedures for salicylate and Nessler's methods. Instructions for conventional laboratory measurement of total ammonia nitrogen by the three methods can be found in a standard water-analysis manual such as *Standard Methods for the Examination of Water and Wastewater*.

(Editor's Note: This article was originally published in the August 2001 print edition of the Global Aquaculture Advocate.)

Author



CLAUDE E. BOYD, PH.D.

Professor, Department of Fisheries and Allied Aquacultures
International Center for Aquaculture and Aquatic Environments
Auburn University
Auburn, AL 36849 USA

boydce1@auburn.edu (<mailto:boydce1@auburn.edu>)

Copyright © 2016–2020 Global Aquaculture Alliance

All rights reserved.